

# A high mountain lizard from Peru: The world's highest-altitude reptile

José Cerdeña<sup>1</sup>, Jackie Farfán<sup>1</sup>, Aarón J. Quiroz<sup>1</sup>

<sup>1</sup> Universidad Nacional de San Agustín de Arequipa, Museo de Historia Natural, Av. Alcides Carrion s/n, Arequipa, Peru

<http://zoobank.org/F5A100CA-B60F-4922-80D7-E2D7634C65FE>

Corresponding author: Jose Cerdeña ([cerdenajoseal@yahoo.es](mailto:cerdenajoseal@yahoo.es))

Academic editor: Johannes Foufopoulos ♦ Received 27 November 2020 ♦ Accepted 4 February 2021 ♦ Published 15 February 2021

## Abstract

Life at high altitudes is particularly challenging for ectothermic animals like reptiles and involves the evolution of specialised adaptations to deal with low temperatures, hypoxia and intense UV radiation. As a result, only very few reptile taxa are able to survive above 5,000 m elevation and herpetological observations from these altitudes are exceedingly rare. We report here an exceptional observation of a lizard population (*Liolaemus* aff. *tacnae*; Reptilia, Squamata) from the high Andes of Peru. During an ascent of Chachani mountain (6,054 m, 16°11'S, 71°32'W), we observed and documented photographically this species living between 5,000 and 5,400 m above sea level. Following a review of literature, we show that this is the highest known record of a reptile species.

## Key Words

*Liolaemus*, *alticolor-bibronii* group, *L. tacnae*, Chachani, Arequipa

## Introduction

Adaptation to high elevation life in reptiles implicates different specialised anatomical and physiological strategies (Huey 1982; He et al. 2013; Reguera et al. 2014; Kouyoumdjian et al. 2019), which scientists are recently beginning to understand at the genetic level (Yang et al. 2014; Yang et al. 2015; Li et al. 2018). Factors identified as important in reptiles for allowing survival at the upper altitudinal limits relate to food availability and capacities for tolerating hypoxia and low temperatures (Navas 2002; Lu et al. 2018; Gangloff et al. 2019). Prior to this publication, the highest known altitude records for reptiles came from species in two lizard genera, *Phrynocephalus* (up to 5,300 m) from the Tibetan plateau and *Liolaemus* (up to 5,176 m) from the Andes (Zhao et al. 1999; Aparicio and Ocampo 2010; Li et al. 2017).

The South American genus *Liolaemus* constitutes one of the most diverse genera of reptiles known on Earth, with more than 270 species described to date (Abdala et al. 2020; Huamaní-Valderrama et al. 2020); it is distributed

from the Central Andes in Peru to Patagonia in Argentina (Pincheira-Donoso et al. 2008; Aguilar et al. 2013; Abdala et al. 2020). This includes the coastal regions of southern Peru and northern Chile (Valladares et al. 2004; Villegas-Paredes et al. 2020), as well as southern Brazil (Verrastro et al. 2003) and demonstrates a broad adaptive radiation of the genus in the region (Abdala et al. 2020). In Peru, 28 species in the genus *Liolaemus* have been described to date (Chaparro et al. 2020; Huamaní-Valderrama et al. 2020; Arapa-Aquino et al. 2021), although recent phylogenetic analyses have shown that this number is likely an underestimate, given the existence of several undescribed lineages or “candidate species” (Aguilar et al. 2017, 2018; Abdala et al. 2020; Huamaní-Valderrama et al. 2020).

## Observations

On 18 October 2020, we ascended the summit of the Chachani mountain (6,054 m), located 20 km north




of Arequipa in south-western Peru. At approximately 09:00 hrs (local time) and at an elevation of 5,000 m, we observed from a distance of approximately five metres two small lizards moving and then hiding between large rocks, making further observations difficult. Later, during the same day, between 10:00 and 12:00 hrs (local time), a total of ten lizards of the same characteristics were observed between 5,000 and 5,400 m of elevation (Fig. 1A), where they were moving amongst the rocks. One individual was photographed at 5,400 m elevation (Fig. 1B), with coordinates 16°10'57"S, 71°32'02"W (Fig. 2). The habitat at the site is characterised by large rocks on the mountain slopes and the almost complete absence of vegetation (Fig. 1A), with only small herbaceous plants growing near the protected underside of rocks. The lizards were exclusively saxicolous, being observed moving only on rocky substrate.

Based on the pattern, colouration and size (approx. 11 cm total length) of the observed animals, we consider them to belong to the *alticolor-bibronii* group of the genus *Liolaemus*. We provisionally assign this population to *Liolaemus* aff. *tacnae*, because of its similarity to the *L. tacnae* which inhabits Chiguata (4,000 m elevation), approx. 30 km SE of Chachani mountain (Aguilar et al. 2013). However, a more detailed analysis is needed, including use of morphological and/or molecular evidence, to determine with more confidence the taxonomic status of this high mountain population.



**Figure 1.** The highest-altitude record for a reptile. **A.** View of the new record location, on the route of ascent from 5,000 m to the summit of Chachani mountain, south-western Peru. **B.** Adult specimen of *L. aff. tacnae* photographed at 5,400 m altitude (place of photography indicated with a red arrow in Fig. 1A).

The present observations represent the highest documented record for the genus *Liolaemus*, surpassing by more than 200 m elevation the previous record holders:



**Figure 2.** Map of the locations of highest altitude records of reptiles listed in Table 1.



**Table 1.** The world’s highest altitude records for three major groups of terrestrial reptiles.

Group	Family	Specie	Altitude (m)	Region	Reference
Lizards	Liolaemidae	<i>Liolaemus</i> aff. <i>tacnae</i>	5000–5400	South America	Present study
	Agamidae	<i>Phrynocephalus erythrurus</i> Zugmayer, 1909	4500–5300	Asia	Zhao et al. 1999; Li et al. 2017
	Liolaemidae	<i>Liolaemus</i> sp.	5060–5176	South America	Aparicio and Ocampo 2010
	Agamidae	<i>Phrynocephalus theobaldi</i> Blyth, 1863	3600–5100	Asia	Zhao et al. 1999; Jin et al. 2017
	Liolaemidae	<i>Liolaemus polystictus</i> Laurent 1992	4043–5016	South America	Olivera and Aguilar 2020
Snakes	Colubridae	<i>Thermophis baileyi</i> Wall, 1907	3600–4900	Asia	Dorge et al. 2007
	Viperidae	<i>Gloydius himalayanus</i> (Gunther, 1864)	1500–4850	Asia	Whitaker and Captain 2004; Saikia et al. 2007
	Dipsadidae	<i>Tachymenis peruviana</i> Wiegmann, 1834	1800–4600	South America	Miranda et al. 2015; Gallardo et al. 2019
Turtles	Testudinidae	<i>Testudo graeca</i> Linnaeus, 1758	0–2090	North Africa	Anadón et al. 2012
	Testudinidae	<i>Gopherus flavomarginatus</i> Legler, 1959	1000–2000	North America	Becerra-López et al. 2017
	Emydidae	<i>Terrapene carolina</i> Linnaeus, 1758	300–1875	North America	Diggins et al. 2016

two Bolivian populations of *Liolaemus* reported between 5,060 and 5,176 m (Aparicio and Ocampo 2010). This population of *L.* aff. *tacnae* represents the reptile species that lives at the highest altitude on Earth (Table 1), exceeding by 100 m the record documented for the Asian species *Phrynocephalus erythrurus*, which inhabits the Tibetan plateau between 4,500 and 5,300 metres elevation (Zhao et al. 1999; Li et al. 2017).

The ongoing global warming trend has probably facilitated the survival of *L.* aff. *tacnae* at this record high site. The genus *Liolaemus* is well-adapted to a broad range of environments (Pincheira-Donoso et al. 2015) and climate change is expected to favour the expansion of species at their upper altitudinal limits (e.g. Wilson et al. 2005, but also see Parmesan 2006 for exceptions). This is particularly true for ectothermic animals, such as the lizards, whose vital functions are known to be directly dependent on ambient temperatures (Sinervo et al. 2018).

Conclusions

We hope that this report will encourage future research into the physiological mechanisms that allow these animals to live at such extremely high elevations through adaptations to low temperatures and low oxygen availability (e.g. Bennett and Ruben 1975; Lu et al. 2015; Cordero et al. 2017; Gangloff et al. 2019).

Acknowledgements

The authors are very grateful to Ivan Jimenez and Ander Tejada for their support in the ascent of the Chachani mountain. To Mr. Libio Roy Santa Cruz of the Herpetology Area of the Natural History Museum of Universidad Nacional de San Agustín de Arequipa (UNSA) for facilitating access to the collections under their care. We also thank Juan Carlos Chaparro for assistance in the preparation of the map. We acknowledge the Subject Editor, Johannes Foufopoulos, for his reading and corrections. Our thanks also go to the reviewers of the document for their valuable comments.

References

Abdala CS, Quinteros AS, Semhan RV, Bulacios-Arroyo AL, Schulte J, Paz MM, Ruiz-Monachesi MR, Laspiur A, Aguilar-Kirigin AJ, Gutiérrez Poblete R, Valladares Faundez P, Valdés J, Portelli S, Santa-Cruz R, Aparicio J, García N, Langstroth R (2020) Unravelling interspecific relationships among highland lizards: first phylogenetic hypothesis using total evidence of the *Liolaemus montanus* group (Iguania: Liolaemidae). *Zoological Journal of the Linnean Society* 189(1): 349–377. <https://doi.org/10.1093/zoolinnean/zlz114>

Aguilar C, Wood Jr PL, Belk M, Duff MH, Sites Jr JW (2017) Different roads lead to Rome: Integrative taxonomic approaches lead to the discovery of two new lizard lineages in the *Liolaemus montanus* group (Squamata: Liolaemidae). *Biological Journal of the Linnean Society* 120: 448–467. <https://doi.org/10.1111/bij.12890>

Aguilar C, Avila LJ, De la Riva I, Johnson L, Morando M, Troncoso-Palacios J, Wood Jr PL, Sites Jr JW (2018) The shadow of the past: convergence of young and old South American desert lizards as measured by head shape traits. *Ecology and Evolution* 8: 11399–11409. <https://doi.org/10.1002/ece3.4548>

Aguilar C, Wood Jr PL, Cusi JC, Guzmán A, Huari F, Lundberg M, Mortensen E, Ramírez C, Robles D, Suárez J, Ticona, A, Vargas VJ, Venegas PJ, Sites Jr JW (2013) Integrative taxonomy and preliminary assessment of species limits in the *Liolaemus walker* complex (Squamata, Liolaemidae) with descriptions of three new species from Peru. *ZooKeys* 364: 47–91. <https://doi.org/10.3897/zookeys.364.6109>

Anadón JD, Giménez A, Graciá E, Pérez I, Rouag R, Larbes S, Kalbousi M, Fahd S, Znari M, El Mouden H, Slimani T, Jdeidi T (2012) Distribution of *Testudo graeca* in Western Mediterranean according to climatic factors. *Amphibia-Reptilia* 33: 285–296. <https://doi.org/10.1163/156853812X643710>

Aparicio J, Ocampo M (2010) *Liolaemus* grupo *montanus* Etheridge, 1995 (Iguania–Liolaemidae). *Novedad zoogeográfica. Cuadernos de Herpetología* 24(2): 133–135. <http://ppct.caicyt.gov.ar/index.php/cuadherpetol/article/view/996/906>

Arapa-Aquino LP, Abdala CS, Huamaní-Valderrama L, Gutierrez RC, Cerdeña JA, Quiroz JA, Chaparro JC (2021) Una nueva especie de lagartija del género *Liolaemus* (Iguania: Liolaemidae) endémica de la Puna del sur de Perú. *Cuadernos de Herpetología*, 35 (Suppl. 1): 35–48. <http://ppct.caicyt.gov.ar/index.php/cuadherpetol/article/view/19508/45454575771299>

Becerra-López JL, García-De la Peña C, Romero-Méndez U, Ramírez-Bautista A (2017) Plant cover effect on Bolson tortoise (*Gopherus flavomarginatus* Legler 1959, Testudinidae) burrow use.



- Nature Conservation 17: 57–69. <https://doi.org/10.3897/natureconservation.17.11582>
- Bennett AF, Ruben J (1975) High altitude adaptation and anaerobiosis in sceloporine lizards. *Comparative Biochemistry and Physiology Part A: Physiology* 50(1): 105–108. [https://doi.org/10.1016/S0010-406X\(75\)80209-X](https://doi.org/10.1016/S0010-406X(75)80209-X)
- Chaparro JC, Quiroz AJ, Mamani L, Gutiérrez RC, Condori P, De la Riva I, Herrera-Juárez G, Cerdeña J, Arapa LP, Abdala CS (2020) An endemic new species of Andean lizard of the genus *Liolaemus* from southern Perú (Iguania: Liolaemidae) and its phylogenetic position. *Amphibian & Reptile Conservation* 14(2): 47–63 (e238). [http://amphibian-reptile-conservation.org/pdfs/Volume/Vol\\_14\\_no\\_2/ARC\\_14\\_2\\_\[General\\_Section\]\\_47-63\\_e238.pdf](http://amphibian-reptile-conservation.org/pdfs/Volume/Vol_14_no_2/ARC_14_2_[General_Section]_47-63_e238.pdf)
- Cordero GA, Andersson BA, Souchet J, Micheli G, Noble DWA, Gangloff EJ, Uller T, Aubret F (2017) Physiological plasticity in lizard embryos exposed to high-altitude hypoxia. *Journal of Experimental Zoology Part A: Ecological and Integrative Physiology* 327(7): 423–432. <https://doi.org/10.1002/jez.2115>
- Diggins CA, Higdon SD, Ford WM (2016) High-elevation record for eastern box turtle in the southern Appalachians. *Herpetological Review* 47(3): e454. <https://ssarherps.org/herpetological-review-pdfs/>
- Dorge T, Hofmann S, Wangdwei M0, Duoje L, Solhoy T, Miehe G (2007) The ecological specialist, *Thermophis baileyi* (Wall, 1907) – new records, distribution and biogeographic conclusions. *The Herpetological Bulletin* 101: 8–12. <https://www.thebhs.org/publications/the-herpetological-bulletin/issue-number-101-autumn-2007/3046-hb101-03?format=html>
- Gallardo GA, Nenda SJ, Scrocchi MGJ (2019) *Tachymenis peruviana* Wiegmann, 1834 (Serpentes, Dipsadidae) in Argentina: geographic distribution and a new province record. *Check List* 15(1): 7–12. <https://doi.org/10.15560/15.1.7>
- Gangloff EJ, Sorlin M, Cordero GA, Souchet J, Aubret F (2019) Lizards at the Peak: Physiological plasticity does not maintain performance in lizards transplanted to high altitude. *Physiological and Biochemical Zoology* 92(2): 189–200. <https://doi.org/10.1086/701793>
- He J, Xiu M, Tang X, Yue F, Wang N, Yang S, Chen Q (2013) The different mechanisms of hypoxic acclimatization and adaptation in lizard *Phrynocephalus vlangalii* living on Qinghai-Tibet Plateau. *Journal of Experimental Zoology Part A: Ecological Genetics and Physiology* 319(3): 117–123. <https://doi.org/10.1002/jez.1776>
- Huamani-Valderrama L, Quiroz AJ, Gutiérrez RC, Aguilar-Kirigin A, Huanca-Mamani W, Valladares-Faúndez P, Cerdeña J, Chaparro JC, Santa Cruz R, Abdala CS (2020) Some color in the desert: description of a new species of *Liolaemus* (Iguania: Liolaemidae) from southern Peru, and its conservation status. *Amphibian & Reptile Conservation* 14(3): 1–30(e250). [http://amphibian-reptile-conservation.org/pdfs/Volume/Vol\\_14\\_no\\_3/ARC\\_14\\_3\\_\[Taxonomy\\_Section\]\\_1-30\\_e250.pdf](http://amphibian-reptile-conservation.org/pdfs/Volume/Vol_14_no_3/ARC_14_3_[Taxonomy_Section]_1-30_e250.pdf)
- Huey RB (1982) Temperature, physiology, and ecology of reptiles. In: Gans C, Pough FH (Eds) *Biology of the Reptilia*. Academic Press, New York, 25–91.
- Jin Y, Liu N, Brown R (2017) The geography and timing of genetic divergence in the lizard *Phrynocephalus theobaldi* on the Qinghai-Tibetan plateau. *Scientific Reports* 7: e2281. <https://doi.org/10.1038/s41598-017-02674-4>
- Kouyoumdjian L, Gangloff EJ, Souchet J, Cordero GA, Dupoué A, Aubret F (2019) Transplanting gravid lizards to high elevation alters maternal and embryonic oxygen physiology, but not reproductive success or hatchling phenotype. *Journal of Experimental Biology* 222: jeb206839. <https://doi.org/10.1242/jeb.206839>
- Li XT, Wang Y, Lu SS, Li M, Men SK, Bai YC, Tang XL, Chen Q (2017) The cold hardiness of *Phrynocephalus erythrurus*, the lizard living at highest altitude in the world. *CryoLetters* 38(3): 216–227. [http://www.cryoletters.org/Abstracts/vol\\_38\\_3\\_2017.htm#216](http://www.cryoletters.org/Abstracts/vol_38_3_2017.htm#216)
- Li, JT, Gao YD, Xie L, Deng C, Shi P, Guan ML, Huang S, Ren JL, Wu DD, Ding L, Huang ZY, Nie H, Humphreys DP, Hillis DM, Wang WZ, Zhang YP (2018) Comparative genomic investigation of high-elevation adaptation in ectothermic snakes. *Proceedings of the National Academy of Sciences of the United States of America* 115: 8406–8411. <https://doi.org/10.1073/pnas.1805348115>
- Lu S, Xin Y, Tang S, Yue F, Wang H, Bai Y, Niu Y, Chen Q (2015) Differences in hematological traits between high- and low-altitude lizards (genus *Phrynocephalus*). *PLoS ONE* 10: e0125751. <https://doi.org/10.1371/journal.pone.0125751>
- Lu HL, Xu CX, Zeng ZG, Du WG (2018) Environmental causes of between-population difference in growth rate of a high-altitude lizard. *BMC Ecology* 18: e37. <https://doi.org/10.1186/s12898-018-0194-8>
- Miranda B, Jimenez RO, Paredes M (2015) Distribución y notas de historia natural de *Tachymenis peruviana* (Wiegmann, 1835) (Dipsadidae: Tachymenini) en Bolivia. *Libro de Resúmenes XVI Congreso Argentino de Herpetología*, San Miguel de Tucumán, Argentina, 306 pp.
- Navas CA (2002) Herpetological diversity along Andean elevational gradients: links with physiological ecology and evolutionary physiology. *Comparative Biochemistry and Physiology-Part A: Molecular & Integrative Physiology* 133: 469–485. [https://doi.org/10.1016/S1095-6433\(02\)00207-6](https://doi.org/10.1016/S1095-6433(02)00207-6)
- Olivera DA, Aguilar CA (2020) Dieta de la lagartija neotropical *Liolaemus polystictus* (Squamata: Liolaemidae) de los andes centrales, Huancavelica, Perú. *Revista peruana de biología* 27(3): 339–348. <https://doi.org/10.15381/rpb.v27i3.18680>
- Parnesan C (2006) Ecological and evolutionary responses to recent climate change. *Annual Review of Ecology, Evolution, and Systematics* 37: 637–669. <https://doi.org/10.1146/annurev.ecolsys.37.091305.110100>
- Pincheira-Donoso D, Sclorano JA, Sura P (2008) A monographic catalogue on the systematics and phylogeny of the South American iguanian lizard family Liolaemidae (Squamata, Iguania). *Zootaxa* 1800: 3–85. <https://doi.org/10.11646/zootaxa.1800.1.1>
- Pincheira-Donoso D, Harvey LP, Ruta M (2015) What defines an adaptive radiation? Macroevolutionary diversification dynamics of an exceptionally species-rich continental lizard radiation. *BMC Evolutionary Biology* 15(1): e153. <https://doi.org/10.1186/s12862-015-0435-9>
- Reguera S, Zamora-Camacho FJ, Moreno-Rueda G (2014) The lizard *Psammmodromus algirus* (Squamata Lacertidae) is darker at high altitudes. *Biological Journal of the Linnean Society* 112: 132–41. <https://doi.org/10.1111/bij.12250>
- Saikia U, Sharma DK, Sharma RM (2007) Checklist of the Reptilian fauna of Himachal Pradesh, India. *Reptile Rap* 8: 6–9. <https://www.zoosprint.zooreach.org/ZoosPrintNewsLetter/ReptileRap%208%20Dec07.pdf>
- Sinervo B, Miles DB, Wu Y, Mendez-de La Cruz F, Kirchhof S, Qi Y (2018) Climate change, thermal niches, extinction risk and maternal-effect rescue of toad-headed lizards, *Phrynocephalus*, in thermal extremes of the Arabian Peninsula to the Qinghai-Tibetan Plateau. *Integrative Zoology* 13: 450–470. <https://doi.org/10.1111/1749-4877.12315>



- Valladares P (2004) Nueva especie de lagarto del género *Liolaemus* (Reptilia: Liolaemidae) del norte de Chile, previamente confundido con *Liolaemus* (= *Phrynosaura*) *reichei*. Cuadernos de Herpetología 18: 41–51.
- Verrastro L, Veronese L, Bujes C, Dias-Filho MM (2003) A new species of *Liolaemus* from Southern Brazil (Iguania: Tropiduridae). Herpetologica 59(1): 105–118. [https://doi.org/10.1655/0018-0831\(2003\)059\[0105:ANSOLF\]2.0.CO;2](https://doi.org/10.1655/0018-0831(2003)059[0105:ANSOLF]2.0.CO;2)
- Villegas-Paredes L, Huamaní-Valderrama L, Luque-Fernández C, Gutiérrez RC, Quiróz AJ, Abdala CS (2020) Una nueva especie de *Liolaemus* (Iguania: Liolaemidae) perteneciente al grupo *L. montanus* en las lomas costeras del sur de Perú. Revista de Biología Tropical 68(1): 69–86. <https://doi.org/10.15517/rbt.v68i1.34861>
- Whitaker R, Captain A (2004) Snakes of India, the Field Guide. Draco Books, Chennai, 481 pp.
- Wilson RJ, Gutiérrez D, Gutiérrez J, Martínez D, Agudo R, Monserrat VJ (2005) Changes to the elevational limits and extent of species ranges associated with climate change. Ecology Letters 8: 1138–1146. <https://doi.org/10.1111/j.1461-0248.2005.00824.x>
- Yang W, Qi Y, Fu J (2014) Exploring the genetic basis of adaptation to high elevations in reptiles: a comparative transcriptome analysis of two toad-headed agamas (genus *Phrynocephalus*). PLoS ONE 9: e112218. <https://doi.org/10.1371/journal.pone.0112218>
- Yang Y, Wang L, Han J, Tang X, Ma M, Wang K, Zhang X, Ren Q, Chen Q, Qiu Q (2015) Comparative transcriptomic analysis revealed adaptation mechanism of *Phrynocephalus erythrurus*, the highest altitude Lizard living in the Qinghai-Tibet Plateau. BMC Evolutionary Biology 15: 101. <https://doi.org/10.1186/s12862-015-0371-8>
- Zhao EM, Zhao KT, Zhou KY (1999) Fauna Sinica: Reptilia (Squamata, Lacertilia) (Vol. 2). Science Press.